

Circular dichroism in assemblies of plasmonic nanoparticles — modelling in the dipole approximation

Baptiste Augu e

28th November 2012

Abstract

The `cda` package implements the coupled-dipole approximation for electromagnetic scattering by sparse collections of subwavelength particles, with a particular focus on plasmonic nanoparticles in the visible regime. The interaction matrix is formed in C++ code for speed; convenient wrapper functions are provided at the R level to calculate the extinction, scattering, and absorption of light by particles with linearly and circularly polarised light. Functions are also provided to calculate orientation-averaged circular dichroism, and display clusters of nanoparticles in three dimensions using RGL or povray.

The system consists in a three-dimensional arrangement of small ellipsoidal particles in arbitrary orientations, as shown in figure 1. Here the position and orientation of the particles was chosen to follow a helix.




Figure 1: (a) Schematic of the 3D arrangement of particles. (b) Orientation of prolate particles in space using two Euler angles.

1 Coupled dipole model

Each dipole scatters light in proportion to the local field it experiences,

$$\mathbf{p}_{\text{dip}} = \alpha \mathbf{E}_{\text{loc}}, \quad (1)$$

where α is the polarizability tensor describing the individual nanoparticle. The prescription of Kuwata *et al.* [1] was chosen, which provides an accurate approximation for particles below ~ 100 nm in size. Each dipole may be placed in arbitrary orientation, and to this end the polarizability tensor must be transformed to a rotated frame.

Let us describe each nanoparticle by a diagonal polarizability tensor in the reference frame of its principal axes. The rotation of this reference frame to the actual position of the particle can be described by three Euler angles ϕ, θ, ψ and a rotation matrix ¹

$$\mathbf{R} = \begin{bmatrix} \cos \psi \cos \phi - \cos \theta \sin \phi \sin \psi & \cos \psi \sin \phi + \cos \theta \cos \phi \sin \psi & \sin \psi \sin \theta \\ -\sin \psi \cos \phi - \cos \theta \sin \phi \cos \psi & -\sin \psi \sin \phi + \cos \theta \cos \phi \cos \psi & \cos \psi \sin \theta \\ \sin \phi \sin \theta & -\cos \phi \sin \theta & \cos \theta \end{bmatrix}. \quad (2)$$

A dipole in orientation ϕ, θ, ψ will be described by a polarizability $\mathbf{R}^{-1}\alpha\mathbf{R}$ in the global reference frame.

The local field

$$\mathbf{E}_{\text{loc}} = \mathbf{E}_{\text{inc}} + \sum_{\text{dipoles} \setminus \text{itself}} \mathbf{E}_{\text{d}}, \quad (3)$$

is the sum of the incident field plus the contribution of the dipolar field associated with the other dipoles in the system. The field radiated by a dipole reads,

$$\mathbf{E}_{\text{d}} = \frac{e^{i\omega r/c}}{4\pi\epsilon_0} \left\{ \frac{\omega^2}{c^2 r} \hat{\mathbf{r}} \times \mathbf{p} \times \hat{\mathbf{r}} + \left(\frac{1}{r^3} - \frac{i\omega}{cr^2} \right) [3(\hat{\mathbf{r}} \cdot \mathbf{p})\hat{\mathbf{r}} - \mathbf{p}] \right\}. \quad (4)$$

By grouping together the dipole moments we can cast equation 3 in matrix form,

$$\mathbf{A}\mathbf{P} = \mathbf{E}_{\text{inc}}, \quad (5)$$

where A is the interaction matrix that describes the electromagnetic coupling between the dipoles in the non-diagonal blocks,

$$A_{ij} = \frac{e^{(ikr_{ij})}}{r_{ij}} \left\{ k^2(\hat{\mathbf{r}} \otimes \hat{\mathbf{r}} - \mathbb{I}) + \frac{ikr_{ij} - 1}{r_{ij}^2} (3\hat{\mathbf{r}} \otimes \hat{\mathbf{r}} - \mathbb{I}) \right\}, \quad (6)$$

and the block diagonal $A_{ii} = \alpha^{-1}$ is formed with the inverse polarizability of the individual dipoles, in the global (x, y, z) reference frame.

When the dipole moments are known by inversion of equation 5, the extinction cross-section can be obtained *for a given incident field* following,

$$\sigma_{\text{ext}} = \frac{4\pi k}{|\mathbf{E}_{\text{inc}}|^2} \Im(\mathbf{E}_{\text{inc}}^* \cdot \mathbf{P}). \quad (7)$$

1.1 Computation of circular dichroism

Circular dichroism can be calculated from the difference in extinction for left-handed and right-handed circularly polarised light, averaged over the full solid angle of incident light,

$$\sigma_{\text{CD}} = \langle \sigma_L \rangle_{\Omega} - \langle \sigma_R \rangle_{\Omega}. \quad (8)$$

The incident field incident along x is written as,

$$\mathbf{E}_{\text{inc}} = \frac{\exp i(\omega t - k_x x)}{\sqrt{2}} \begin{pmatrix} 0 \\ i \\ 1 \end{pmatrix} \quad (\text{right-handed}) \quad (9)$$

$$\mathbf{E}_{\text{inc}} = \frac{\exp i(\omega t - k_x x)}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ i \end{pmatrix} \quad (\text{left-handed}). \quad (10)$$

¹using the same conventions as <http://mathworld.wolfram.com/EulerAngles.html>

More generally, the incident beam will be characterised by a wave-vector \mathbf{k} and an electric vector \mathbf{E}_{inc} describing the light polarisation, and both of these vectors can be rotated using the rotation matrix R as $R^{-1}\mathbf{k}$ and $R^{-1}\mathbf{E}_{\text{inc}}$.

The interaction matrix A does not depend on the direction of the incident field, it may therefore be advantageous to compute A^{-1} (at each wavelength) and perform the matrix-vector products for all the required incident fields. The CD spectra obtained experimentally are averaged over all orientations of the incident beam, it is therefore necessary to use incident wave-vectors that span the full range of $\phi \in [0, 2\pi], \theta = \pi/2, \psi \in [-\pi/2, \pi/2]$.

Averaging the extinction cross-section over all incident field directions is performed by numerical integration,

$$\langle \sigma \rangle_{\Omega} = \frac{1}{4\pi} \int_0^{2\pi} \int_{-\pi/2}^{\pi/2} \sigma(\phi, \psi) \cos \psi d\psi d\phi. \quad (11)$$

A Gauss-Legendre quadrature scheme is used to perform the integration, so that the evaluation of the extinction cross-section is performed for a relatively low number of angles (20×20 seems sufficient). Other integration schemes are also possible.

References

- [1] Hitoshi Kuwata, Hiroharu Tamaru, Kunio Esumi, and Kenjiro Miyano. Resonant light scattering from metal nanoparticles: Practical analysis beyond Rayleigh approximation. 83(22):4625–4627, 2003.
- [2] Baptiste Auguié and William L. Barnes. Diffractive coupling in gold nanoparticle arrays and the effect of disorder. *Opt. Lett.*, 34(4):401–403, Feb 2009.
- [3] Baptiste Auguié, José Lorenzo Alonso-Gómez, Andrés Guerrero-Martínez, and Luis M. Liz-Marzán. Fingers crossed: Optical activity of a chiral dimer of plasmonic nanorods. *The Journal of Physical Chemistry Letters*, 2(8):846–851, 2011.
- [4] Andrés Guerrero-Martínez, Baptiste Auguié, José Lorenzo Alonso-Gómez, Zoran Džolić, Sergio Gómez-Graña, Mladen Žinić, M. Magdalena Cid, and Luis M. Liz-Marzán. Intense optical activity from three-dimensional chiral ordering of plasmonic nanoantennas. *Angewandte Chemie International Edition*, 50(24):5499–5503, 2011.
- [5] Andrés Guerrero-Martínez, José Lorenzo Alonso-Gómez, Baptiste Auguié, M. Magdalena Cid, and Luis M. Liz-Marzán. From individual to collective chirality in metal nanoparticles. *Nano Today*, in press, 2011.
- [6] Baptiste Auguié, Xesús M. Bendaña, William L. Barnes, and F. Javier García de Abajo. Diffractive arrays of gold nanoparticles near an interface: Critical role of the substrate. *Phys. Rev. B*, 82(15):155447, Oct 2010.
- [7] Baptiste Auguié and William Barnes. Collective resonances in gold nanoparticle arrays. 101(14):143902, Sep 2008.