

# Package ‘photobiologyWavebands’

October 24, 2023

**Type** Package

**Title** Waveband Definitions for UV, VIS, and IR Radiation

**Version** 0.5.2

**Date** 2023-10-24

**Maintainer** Pedro J. Aphalo <pedro.aphalo@helsinki.fi>

**Description** Constructors of waveband objects for commonly used biological spectral weighting functions (BSWFs) and for different wavebands describing named ranges of wavelengths in the ultraviolet (UV), visible (VIS) and infrared (IR) regions of the electromagnetic spectrum. Part of the 'r4photobiology' suite, Aphalo P. J. (2015) <[doi:10.19232/uv4pb.2015.1.14](https://doi.org/10.19232/uv4pb.2015.1.14)>.

**License** GPL (>= 2)

**VignetteBuilder** knitr

**Depends** R (>= 4.0.0), photobiology (>= 0.11.0)

**Suggests** knitr (>= 1.40), rmarkdown (>= 2.20)

**LazyLoad** yes

**LazyData** yes

**ByteCompile** true

**Encoding** UTF-8

**URL** <https://docs.r4photobiology.info/photobiologyWavebands/>,  
<https://github.com/aphalo/photobiologywavebands>

**BugReports** <https://github.com/aphalo/photobiologywavebands/issues>

**RoxygenNote** 7.2.3

**NeedsCompilation** no

**Author** Pedro J. Aphalo [aut, cre] (<<https://orcid.org/0000-0003-3385-972X>>),  
Titta K. Kotilainen [ctb] (<<https://orcid.org/0000-0002-2822-9734>>)

**Repository** CRAN

**Date/Publication** 2023-10-24 14:50:03 UTC

**R topics documented:**

photobiologyWavebands-package	3
Blue	5
CH4	7
CH4_e_fun	8
CH4_q_fun	9
CIE1924_1ef.spct	9
CIE1951_scotopic_1ef.spct	10
CIE2008_1ef2deg.spct	11
CIE_e_fun	11
CIE_q_fun	12
DNA_GM	13
DNA_GM_q_fun	13
DNA_N	14
DNA_N_q_fun	15
DNA_P	16
DNA_P_q_fun	16
erythema	17
Far_red	18
FLAV	20
FLAV_q_fun	21
GEN_G	22
GEN_G_q_fun	23
GEN_M	24
GEN_M_q_fun	25
GEN_T	26
GEN_T_q_fun	27
Green	28
ICNIRP_e_fun	29
IR	30
IR_bands	31
Landsat_bands	32
NDVI	33
Orange	34
PAR	35
PG	37
PG_q_fun	39
photopic_sensitivity	40
Plant_bands	40
Purple	41
Red	42
scotopic_sensitivity	44
SetlowTUV.spct	44
UV	45
UVA	46
UVB	47
UVC	49

UVI . . . . .	50
UV_bands . . . . .	51
UV_health_hazard . . . . .	52
VIS . . . . .	53
VIS_bands . . . . .	54
Yellow . . . . .	55
<b>Index</b>	<b>56</b>

---

photobiologyWavebands-package

*photobiologyWavebands: Waveband Definitions for UV, VIS, and IR Radiation*

---

## Description

Constructors of waveband objects for commonly used biological spectral weighting functions (BSWFs) and for different wavebands describing named ranges of wavelengths in the ultraviolet (UV), visible (VIS) and infrared (IR) regions of the electromagnetic spectrum. Part of the 'r4photobiology' suite, Aphalo P. J. (2015) [doi:10.19232/uv4pb.2015.1.14](https://doi.org/10.19232/uv4pb.2015.1.14).

## Details

Package 'photobiologyWavebands' provides constructors for objects of class waveband from package 'photobiology'. These constructors are based on standard definitions or frequently used non-standardized definitions. When different definitions are in common use for a given named waveband the constructors accept an argument to chose among them. Whenever an ISO standard provides a definition, this is used by default. In the infrared (IR) there are many different definitions and waveband names in use. We have tried to include most of the commonly used names and definitions.

Definitions "matching" the different bands of Landsat imagers are included. These are simple wavelength ranges for wavelengths at half-maximum response as given in the NASA literature, which in some cases presents small inconsistencies. These definitions cannot exactly reproduce instrument responses as they do not describe the real spectral responsiveness of the satellite imagers.

By necessity we cover only a subset of all definitions in use. These should be thought as convenience functions, as waveband objects according to any arbitrary definition can be constructed with the constructor provided by package [photobiology-package](#).

## Author(s)

**Maintainer:** Pedro J. Aphalo <[pedro.aphalo@helsinki.fi](mailto:pedro.aphalo@helsinki.fi)> ([ORCID](#))

Other contributors:

- Titta K. Kotilainen <[titta.kotilainen@helsinki.fi](mailto:titta.kotilainen@helsinki.fi)> ([ORCID](#)) [contributor]

## References

- Aphalo, Pedro J. (2015) The r4photobiology suite. *UV4Plants Bulletin*, 2015:1, 21-29. doi:10.19232/uv4pb.2015.1.14.
- Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at <https://hdl.handle.net/10138/37558>
- Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) *Photophysiology*, Academic Press, 1971, 6, 131-177
- Diffey, B. L. 1991. Solar ultraviolet radiation effects on biological systems. *Review in Physics in Medicine and Biology* 36 (3): 299-328.
- Green, A. E. S., Miller, J. H. (1975) Measures of biologically active radiation in the 280-340 nm region. Impacts of climate change on the environment. CIAP Monograph, 5, Part 1, Chapter 2.2.4
- Green, A. E. S.; Sawada, T. & Shettle, E. P. (1974) The middle ultraviolet reaching the ground *Photochemistry and Photobiology*, 1974, 19, 251-259
- Ibdah, M., Krins, A., Seidlitz, H. K., Heller, W., Strack, D. & Vogt, T. (2002) Spectral dependence of flavonol and betacyanin accumulation in *Mesembryanthemum crystallinum* under enhanced ultraviolet radiation. *Plant, Cell & Environment*, 25, 1145-1154
- International Commission on Non-Ionizing Radiation Protection (2004) ICNIRP Guidelines on Limits of Exposure to Ultraviolet Radiation of Wavelengths Between 180 nm and 400 nm (Incoherent Optical Radiation). *Health Physics* 87(2):171-186. <https://www.icnirp.org/cms/upload/publications/ICNIRPUV2004.pdf>
- ISO (2007) Optics and photonics - Spectral bands. ISO Standard 20473:2007. ISO, Geneva.
- ISO (2007) Space environment (natural and artificial) - Process for determining solar irradiances. ISO Standard 21348. ISO, Geneva.
- Quaite, F. E., Sutherland, B. M., Sutherland, J. C. Action spectrum for DNA damage in alfalfa lowers predicted impact of ozone depletion. *Nature*, 1992, 358, 576-578
- Leutner, B. and Horning, N. (2016). RStoolbox: Tools for Remote Sensing Data Analysis. R package version 0.1.6. <https://CRAN.R-project.org/package=RStoolbox>
- Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range *Photochemistry and Photobiology*, 78, 456-461
- Musil, C. F. (1995) Differential effects of elevated ultraviolet-B radiation on the photochemical and reproductive performances of dicotyledonous and monocotyledonous arid-environment ephemerals. *Plant, Cell and Environment*, 18, 844-854
- Murakami, K., Aiga I. (1994) Red/Far-red photon flux ratio used as an index number for morphological control of plant growth under artificial lighting conditions. *Proc. Int. Symp. Artificial Lighting, Acta Horticulturae*, 418, ISHS 1997.
- NASA (nd) Landsat 7 Science Data Users Handbook. [https://landsat.gsfc.nasa.gov/wp-content/uploads/2016/08/Landsat7\\_Handbook.pdf](https://landsat.gsfc.nasa.gov/wp-content/uploads/2016/08/Landsat7_Handbook.pdf) Visited on 2016-12-26.
- Sellaro, R., Crepy, M., Trupkin, S. A., Karayekov, E., Buchovsky, A. S., Rossi, C., & Casal, J. J. (2010). Cryptochrome as a sensor of the blue/green ratio of natural radiation in *Arabidopsis*. *Plant physiology*, 154(1), 401-409. doi:10.1104/pp.110.160820

Setlow, R. B. (1974) The Wavelengths in Sunlight Effective in Producing Skin Cancer: A Theoretical Analysis. Proceedings of the National Academy of Sciences, 71, 3363-3366

Smith, H. (1982) Light quality, photoperception and plant strategy. Annual Review of Plant Physiology, 33:481-518.

USGS (nd) Landsat 8 Science Data Users Handbook. <https://www.usgs.gov/media/files/landsat-8-data-users-handbook>. Visited on 2023-01-07.

Webb, A. R.; Slaper, H.; Koepke, P. & Schmalwieser, A. W. Know your standard: clarifying the CIE erythema action spectrum. Photochemistry and photobiology, 2011, 87, 483-486

### See Also

Useful links:

- <https://docs.r4photobiology.info/photobiologyWavebands/>
- <https://github.com/aphalo/photobiologywavebands>
- Report bugs at <https://github.com/aphalo/photobiologywavebands/issues>

### Examples

```
q_irrad(sun.spct, PAR()) # PAR photon irradiance
q_irrad(sun.spct, Blue("ISO")) # blue photon irradiance, ISO definition
q_irrad(sun.spct, Blue("Sellaro")) # blue photon irradiance, Sellaro et al.'s definition
e_irrad(sun.spct, VIS()) # VIS irradiance, ISO definition
q_irrad(sun.spct, VIS()) # VIS photon, ISO definition
```

---

Blue

*Constructor of blue waveband*

---

### Description

Wavelength-range definitions for *blue* light according to ISO or as commonly used in plant or remote sensing applications.

### Usage

```
Blue(std = "ISO")
```

### Arguments

`std` a character string "ISO", "Sellaro", "Broad", "RS" (remote sensing), or Landsat imagers, "LandsatTM", "LandsatETM", or "LandsatOLI".

**Details**

The different arguments passed to formal parameter `std` determine the range of wavelengths set as boundaries of the returned waveband object; "ISO" is standardized definition based on human colour vision; "Sellaro" and "Broad" are non-standard but used in plant sciences; "RS" is non-standard but frequently used in remote sensing; the remaining definitions are for the published wavelength sensitivity range of imagers (cameras) in the Landsat satellite missions.

**Value**

A waveband object defining a wavelength range.

**Note**

The bands are defined as square windows, these can be applied to spectral data to obtain the "true" values, but they do not simulate the sensitivity of broad-band sensors or the spectral transmittance of ionic filters. Some band-pass interference filters may have very sharp cut-in and cut-off, and their effect can be approximated by a square window, but filters based on light absorption will show gradual tails and bell-shaped wavelength-windows. The Landsat instruments have very steep cut-in and cut-off slopes and are well approximated.

**References**

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

ISO (2007) *Space environment (natural and artificial) - Process for determining solar irradiances*. ISO Standard 21348. ISO, Geneva.

Sellaro, R., Crepy, M., Trupkin, S. A., Karayekov, E., Buchovsky, A. S., Rossi, C., & Casal, J. J. (2010). Cryptochrome as a sensor of the blue/green ratio of natural radiation in Arabidopsis. *Plant physiology*, 154(1), 401-409. [doi:10.1104/pp.110.160820](https://doi.org/10.1104/pp.110.160820).

**See Also**

[new\\_waveband](#) [waveband](#)

Other unweighted wavebands: [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

**Examples**

```
Blue()
Blue("ISO")
Blue("Sellaro")

e_irrad(sun.spct, Blue()) # W m-2
q_irrad(sun.spct, Blue()) # mol m-2
q_irrad(sun.spct, Blue(), scale.factor = 1e6) # umol m-2
```

---

CH4

*Constructor of CH4 production from pectin weighted waveband*

---

### Description

Methane production from pectin BSWF

### Usage

```
CH4(norm = 300, w.low = 275, w.high = 400)
```

### Arguments

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

### Value

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

### References

Bloom, A. A.; Lee-Taylor, J.; Madronich, S.; Messenger, D. J.; Palmer, P. I.; Reay, D. S. & McLeod, A. R. (2010) Global methane emission estimates from ultraviolet irradiation of terrestrial plant foliage. *New Phytologist*, Blackwell Publishing Ltd, 187, 417–425 .

### See Also

[waveband](#)

Other BSWF weighted wavebands: [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

### Examples

```
CH4()  
CH4(norm = 400)
```

---

CH4_e_fun	<i>Gives values for the CH4 production from pectin BSWF as a function of wavelength</i>
-----------	---

---

### Description

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on quantum based effectiveness relative units.

### Usage

```
CH4_e_fun(w.length)
```

### Arguments

w.length          numeric array of wavelengths (nm)

### Value

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (300 nm) and based on energy effectiveness.

### References

Bloom, A. A.; Lee-Taylor, J.; Madronich, S.; Messenger, D. J.; Palmer, P. I.; Reay, D. S. & McLeod, A. R. (2010) Global methane emission estimates from ultraviolet irradiation of terrestrial plant foliage. *New Phytologist*, Blackwell Publishing Ltd, 187, 417–425 .

### See Also

Other BSWF functions: [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

### Examples

```
CH4_e_fun(293:400)
```



---

CH4_q_fun	<i>Gives values for the CH4 production from pectin BSWF as a function of wavelength</i>
-----------	---

---

### Description

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on quantum based effectiveness relative units.

### Usage

```
CH4_q_fun(w.length)
```

### Arguments

w.length          numeric array of wavelengths (nm)

### Value

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (300 nm) but based on quantum effectiveness.

### See Also

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

### Examples

```
CH4_q_fun(293:400)
```

---

CIE1924_1ef.spct	<i>CIE1924 luminous efficiency function (photopic human vision)</i>
------------------	---

---

### Description

A dataset containing the wavelengths at a 1 nm interval. Tabulated values for quantum luminous efficiency according to CIE1924.

### Format

A response.spct object with 471 rows and 2 variables

**Details**

The variables are as follows:

- w.length (nm)
- s.q.response

**Note**

This luminous efficiency function underestimates the response to short wavelengths.

**References**

<http://www.cvr1.org/> downloaded on 2015-01-24

---

CIE1951\_scotopic\_lef.spct

*Luminous efficiency function (scotopic human vision)*

---

**Description**

A dataset containing the wavelengths at a 1 nm interval. Tabulated values for quantum luminous efficiency at low light levels according to CIE1951.

**Format**

A response.spct object with 401 rows and 2 variables

**Details**

The variables are as follows:

- w.length (nm)
- s.q.response

**References**

<http://www.cvr1.org/> downloaded on 2015-01-24

---

CIE2008\_lef2deg.spct    *CIE2008 luminous efficiency function (2-deg) (photopic human vision)*

---

**Description**

A dataset containing the wavelengths at a 1 nm interval. Tabulated values for quantum luminous efficiency according to CIE2008 for 2 degrees.

**Format**

A response.spct object with 441 rows and 2 variables

**Details**

The variables are as follows:

- w.length (nm)
- s.q.response

**References**

<http://www.cvr1.org/> downloaded on 2015-01-24

---

CIE\_e\_fun                      *Gives values for the erythema BSWF as a function of wavelength*

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on quantum based effectiveness relative units.

**Usage**

```
CIE_e_fun(w.length)
```

**Arguments**

w.length                      numeric array of wavelengths (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (298 nm) and based on energy effectiveness.

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
CIE_e_fun(293:400)
```

---

CIE\_q\_fun

*Gives values for the erythemal BSWF as a function of wavelength*

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on quantum based effectiveness relative units.

**Usage**

```
CIE_q_fun(w.length)
```

**Arguments**

w.length          numeric array of wavelengths (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (298 nm) and based on quantum effectiveness.

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
CIE_q_fun(293:400)
```

---

DNA_GM	<i>Constructor of DNA damage (SETLOW) weighted waveband</i>
--------	---

---

**Description**

Naked DNA damage BSWF, Green and Miller's formulation.

**Usage**

```
DNA_GM(norm = 300, w.low = 275, w.high = 400)
```

**Arguments**

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**See Also**

[new\\_waveband waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
DNA_GM()
DNA_GM(300)
```

---

DNA_GM_q_fun	<i>Gives values for naked DNA BSWF (SETLOW) as a function of wavelength</i>
--------------	---

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. It uses the seldom used Green and Miller formulation.

**Usage**

```
DNA_GM_q_fun(w.length)
```

**Arguments**

w.length          numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
DNA_GM_q_fun(293:400)
```

---

DNA\_N

*Constructor of DNA damage (SETLOW) weighted waveband*

---

**Description**

Naked DNA damage BSWF

**Usage**

```
DNA_N(norm = 300, w.low = 275, w.high = 400)
```

**Arguments**

norm                normalization wavelength (nm)  
w.low                short-end boundary wavelength (nm)  
w.high                long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**See Also**

[new\\_waveband](#) [waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
DNA_N()  
DNA_N(300)
```

---

DNA_N_q_fun	<i>Gives values for naked DNA BSWF (SETLOW) as a function of wavelength</i>
-------------	---

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances.

**Usage**

```
DNA_N_q_fun(w.length)
```

**Arguments**

w.length          numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**Note**

The digitized data as used in the TUV model covers the wavelength range from 256 nm to 364 nm. For longer wavelengths we set the value to zero, and for shorter wavelengths we extrapolate the value for 256 nm.

**Examples**

```
DNA_N_q_fun(293:400)
```

---

DNA_P	<i>Constructor of DNA damage (Quaite) weighted waveband</i>
-------	---

---

**Description**

Plant DNA damage BSWF as formulated by Musil.

**Usage**

```
DNA_P(norm = 300, w.low = 275, w.high = 400)
```

**Arguments**

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**See Also**

[new\\_waveband waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
DNA_P()
DNA_P(300)
```

---

DNA_P_q_fun	<i>Gives values for plant DNA BSWF (Quaite) as a function of wavelength</i>
-------------	---

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. It uses the formulation proposed by Musil.

**Usage**

```
DNA_P_q_fun(w.length)
```



**Arguments**

w.length          numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
DNA_P_q_fun(293:400)
```

---

erythema

*Constructor of erythema-weighted waveband*

---

**Description**

Erythema BSWF (1998 update)

**Usage**

```
erythema(std = "CIE98", norm = 298, w.low = 250, w.high = 400)
```

```
CIE(norm = 298, w.low = 250, w.high = 400)
```

**Arguments**

std                  a character string, currently only "CIE98" supported.

norm                normalization wavelength (nm)

w.low                short-end boundary wavelength (nm)

w.high               long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**Note**

The erythema BSWF from CIE is specified by a mathematical formula, and this is used directly in the definition of the returned waveband.

Standard DIN 5031-10:2018-03 defines BSWF *er* as a table of interpolated values derived from CIE's definition from 1998. So, the values computed using this R package do not necessarily exactly match those according to DIN 5031-10:2018-03. The range of wavelengths used, 250 to 400 nm, does agree, with those in the standard.

**References**

Webb, A. R.; Slaper, H.; Koepke, P. & Schmalwieser, A. W. (2011) Know your standard: clarifying the CIE erythema action spectrum. *Photochemistry and photobiology*, 2011, 87, 483-486

DIN (2018) Standard DIN 5031-10:2018-03 Optical radiation physics and illuminating engineering. Part 10: Photobiologically effective radiation, quantities, symbols and action spectra. Beuth Verlag, Berlin 2018.

**See Also**

[waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#)

**Examples**

```
erythema()
erythema("CIE98")
CIE()
CIE(norm = 300)
erythema(norm = 300)
```

---

Far\_red

*Constructor of far-red waveband*

---

**Description**

Wavelength-range definitions for *far-red* light according as commonly used in plant or remote sensing applications.

**Usage**

```
Far_red(std = "ISO")
```

## Arguments

`std` a character string, defaults to "ISO", as for other colour definitions, which in this case returns NA.

## Details

The different arguments passed to formal parameter `std` determine the range of wavelengths set as boundaries of the returned waveband object; far-red is not defined by "ISO" standard definitions based on human colour vision, and included under red; "Smith10", "Smith20", "Inada", "Warrington", "Sellaro" and "Broad" are non-standard but used in plant sciences; "RedEdge40" and "RedEdge40" are non-standard but frequently used in remote sensing.

In plant photobiology the definitions proposed by Prof. Harry Smith are the most widely used, specially to compute a red to far-red photon ratio relevant to phytochrome photoreceptors. However, other authors have used different definitions in their publications. "Smith10" (725-735 nm), "Smith20" (720-740 nm), "Inada" (700-800 nm), "Warrington" (700-850 nm), and "Sellaro" (700-750 nm).

Other definitions used in remote sensing. For example the "red edge" is to detect the condition of vegetation based on light reflectance by green vegetation. These bands are centred at the reflectance transition in the far-red band (725 nm), and here we define "RedEdge40" (705-745 nm) and "RedEdge20" (715-735 nm).

## Value

A waveband object defining a wavelength range.

## Note

The bands are defined as square windows, these can be applied to spectral data to obtain the "true" values, but they do not simulate the sensitivity of broad-band sensors or the spectral transmittance of ionic filters. Some band-pass interference filters may have very sharp cut-in and cut-off, and their effect can be approximated by a square window, but filters based on light absorption will show gradual tails and bell-shaped wavelength-windows.

## References

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

ISO (2007) Space environment (natural and artificial) - Process for determining solar irradiances. ISO Standard 21348. ISO, Geneva.

Murakami, K., Aiga I. (1994) Red/Far-red photon flux ratio used as an index number for morphological control of plant growth under artificial lighting conditions. Proc. Int. Symp. Artificial Lighting, Acta Horticulturae, 418, ISHS 1997.

Sellaro, R., Crepy, M., Trupkin, S. A., Karayekov, E., Buchovsky, A. S., Rossi, C., & Casal, J. J. (2010). Cryptochrome as a sensor of the blue/green ratio of natural radiation in Arabidopsis. *Plant physiology*, 154(1), 401-409. [doi:10.1104/pp.110.160820](https://doi.org/10.1104/pp.110.160820).

Smith, H. (1982) Light quality, photoperception and plant strategy. Annual Review of Plant Physiology, 33:481-518. doi:10.1146/annurev.pp.33.060182.002405

### See Also

NIR for wavebands close to the boundary between red and infrared regions.

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

### Examples

```
Far_red() # no ISO definition exists
Far_red("ISO") # no ISO definition exists
Far_red("Smith10") # 10 nm wide
Far_red("Smith20") # 20 nm wide
Far_red("Inada")
Far_red("Warrington")
Far_red("Sellarò")
Far_red("RedEdge40")
Far_red("RedEdge20")

e_irrad(sun.spct, Far_red("Smith10")) # W m-2
q_irrad(sun.spct, Far_red("Smith10")) # mol m-2
q_irrad(sun.spct, Far_red("Smith10"), scale.factor = 1e6) # umol m-2
```

---

FLAV

*Constructor of FLAV BSWF flavonoids*

---

### Description

Mesembryanthin accumulation BSWF, data and formulation from Ibdah et al.

### Usage

```
FLAV(norm = 300, w.low = 275, w.high = 346)
```

### Arguments

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

### Value

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

## References

Ibdah, M.; Krins, A.; Seidlitz, H. K.; Heller, W.; Strack, D. & Vogt, T. (2002) Spectral dependence of flavonol and betacyanin accumulation in *Mesembryanthemum crystallinum* under enhanced ultraviolet radiation. *Plant, Cell & Environment*, 25, 1145-1154. doi:10.1046/j.1365-3040.2002.00895.x

## See Also

[waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

## Examples

```
FLAV()
FLAV(300)
```

---

FLAV_q_fun	<i>Gives values for FLAV BSWF (flavonoid) as a function of wavelength</i>
------------	---

---

## Description

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. It is the action spectrum for the accumulation of mesembryanthin.

## Usage

```
FLAV_q_fun(w.length)
```

## Arguments

w.length          numeric array of w.length (nm)

## Value

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

## See Also

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

## Examples

```
FLAV_q_fun(293:400)
```

---

GEN\_G                                      *Constructor of GPAS (Green's formulation) weighted waveband*

---

### Description

Generalized Plant Action BSWF of Caldwell as formulated by Green et al.

### Usage

```
GEN_G(norm = 300, w.low = 275, w.high = 313.3)
```

### Arguments

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

### Value

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

### Note

In the original publication [2] describing the formulation, the long-end wavelength boundary is specified as 313.3 nm. This is the default used here. However, in some cases it is of interest to vary this limit in sensitivity analyses. The effect on the RAF and doses of changing this boundary is substantial, and has been analysed by Micheletti et al. [3].

### References

- [1]Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) Photophysiology, Academic Press, 1971, 6, 131-177
- [2] Green, A. E. S.; Sawada, T. & Shettle, E. P. (1974) The middle ultraviolet reaching the ground Photochemistry and Photobiology, 1974, 19, 251-259
- [3] Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range Photochemistry and Photobiology, 78, 456-461

### See Also

[waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
GEN_G()  
GEN_G(300)
```

---

GEN_G_q_fun	<i>Gives values for GPAS BSWF (Green's formulation) as a function of wavelength</i>
-------------	---

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The BSWF is normalized at 280 nm.

**Usage**

```
GEN_G_q_fun(w.length)
```

**Arguments**

w.length          numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**Note**

In the original publication [2] describing the formulation, the long-end wavelength boundary is specified as 313.3 nm. The equation is coded here with no such limit so that any limit can be set when defining the waveband. We do so because in some cases it is of interest to vary this limit in sensitivity analyses. The effect on the RAF and doses of changing this boundary is substantial, and has been analysed by Micheletti et al. [3].

**References**

- [1] Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) *Photophysiology*, Academic Press, 1971, 6, 131-177
- [2] Green, A. E. S.; Sawada, T. & Shettle, E. P. (1974) The middle ultraviolet reaching the ground *Photochemistry and Photobiology*, 1974, 19, 251-259
- [3] Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range *Photochemistry and Photobiology*, 78, 456-461

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
GEN_G_q_fun(293:400)
```

---

 GEN\_M

---

*Constructor of GPAS (Micheletti's formulation) weighted waveband*


---

**Description**

Generalized Plant Action BSWF of Caldwell [1] as formulated by Micheletti et al. [2]

**Usage**

```
GEN_M(norm = 300, w.low = 275, w.high = 313.3)
```

**Arguments**

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**Note**

In the original publication [2] describing the formulation, the long-end wavelength boundary is specified as 313.3 nm. This is the default used here. However, in some cases it is of interest to vary this limit in sensitivity analyses. The effect on the RAF and doses of changing this boundary is substantial, and has been analysed by Micheletti et al. [3].

**References**

- [1]Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) *Photophysiology*, Academic Press, 1971, 6, 131-177
- [2] Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range *Photochemistry and Photobiology*, 78, 456-461



**See Also**

[new\\_waveband](#) and [waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
GEN_M()
GEN_M(300)
```

---

GEN_M_q_fun	<i>Gives values for GPAS BSWF (Micheletti's formulation) as a function of wavelength</i>
-------------	--

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The BSWF is normalized at 300 nm.

**Usage**

```
GEN_M_q_fun(w.length)
```

**Arguments**

w.length            numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**Note**

In the original publication [2] describing the formulation, the long-end wavelength boundary is not specified, but 313.3 nm is usually used. The equation is coded here with the limit at 342 nm as at longer wavelengths the values increase with increasing wavelength. The effect on the RAF and doses of changing this boundary can be substantial, and has been analysed by Micheletti et al. [3].

**References**

- [1]Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) Photophysiology, Academic Press, 1971, 6, 131-177
- [2] Micheletti, M. I. and R. D. Piacentini (2002) Irradiancia espectral solar UV-B y su relación con la efectividad de daño biológico a las plantas. ANALES AFA, 13, 242-248
- [3] Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range Photochemistry and Photobiology, 78, 456-461

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
GEN_M_q_fun(293:400)
```

---

 GEN\_T

---

*Constructor of GPAS (Thimijan's formulation) weighted waveband*


---

**Description**

Generalized Plant Action BSWF of Caldwell [1] as formulated by Timijan et al. [2]

**Usage**

```
GEN_T(norm = 300, w.low = 275, w.high = 345)
```

**Arguments**

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

**Value**

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

**References**

- [1] Caldwell, M. M. (1971) Solar UV irradiation and the growth and development of higher plants. In Giese, A. C. (Ed.) *Photophysiology*, Academic Press, 1971, 6, 131-177
- [2] Thimijan RW, Cams HR, Campbell L. (1978) Radiation sources and related environmental control for biological and climatic effects of UV research. Final report EPA-IAG-D6-0168. Washington: Environmental Protection Agency.

**See Also**

[GEN.G](#) [GEN.M](#) [PG](#) and [waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [PAR\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

**Examples**

```
GEN_T()  
GEN_T(300)
```

---

GEN_T_q_fun	<i>Gives values for GPAS BSWF (Timijan's formulation) as a function of wavelength</i>
-------------	---

---

**Description**

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances.

**Usage**

```
GEN_T_q_fun(w.length)
```

**Arguments**

w.length          numeric array of w.length (nm)

**Value**

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source. The returned values are based on quantum effectiveness units.

**Note**

For wavelengths shorter than 256 nm the value returned by the equation starts decreasing, but we instead extrapolate this maximum value, obtained at 256 nm, to shorter wavelengths. For wavelengths longer than 345 nm we return zero, as is usual practice.

**See Also**

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

**Examples**

```
GEN_T_q_fun(293:400)
```

---

Green

*Constructor of green waveband*

---

### Description

Wavelength-range definitions for *green* light according to ISO or as commonly used in plant or remote sensing applications.

### Usage

```
Green(std = "ISO")
```

### Arguments

`std` a character string "ISO", "Sellaro", "Broad", "RS", "LandsatOLI", "LandsatRBV", "LandsatTM", "LandsatETM", or "LandsatMSS".

### Details

The different arguments passed to formal parameter `std` determine the range of wavelengths set as boundaries of the returned waveband object; "ISO" is an standardized definition based on human colour vision; "Sellaro" and "Broad" are non-standard but used in plant sciences; "RS" is non-standard but frequently used in remote sensing; the remaining definitions are for the published wavelength sensitivity range of imagers (cameras) in the Landsat satellite missions.

### Value

A waveband object defining a wavelength range.

### Note

The bands are defined as square windows, these can be applied to spectral data to obtain the "true" values, but they do not simulate the sensitivity of broad-band sensors or the spectral transmittance of ionic filters. Some band-pass interference filters may have very sharp cut-in and cut-off, and their effect can be approximated by a square window, but filters based on light absorption will show gradual tails and bell-shaped wavelength-windows. The Landsat instruments have very steep cut-in and cut-off slopes and are well approximated.

When released, this package will replace the package UVcalc.

### References

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

ISO (2007) Space environment (natural and artificial) - Process for determining solar irradiances. ISO Standard 21348. ISO, Geneva.

Sellaro, R., Crepy, M., Trupkin, S. A., Karayekov, E., Buchovsky, A. S., Rossi, C., & Casal, J. J. (2010). Cryptochrome as a sensor of the blue/green ratio of natural radiation in Arabidopsis. *Plant physiology*, 154(1), 401-409. doi:10.1104/pp.110.160820.

### See Also

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

### Examples

```
Green()
Green("ISO") # 500 to 570
Green("Sellaro") # 500 to 570 nm

e_irrad(sun.spct, Green()) # W m-2
q_irrad(sun.spct, Green()) # mol m-2
q_irrad(sun.spct, Green(), scale.factor = 1e6) # umol m-2
```

---

ICNIRP\_e\_fun

*ICNIRP UV health hazard BSWF as a function of wavelength*

---

### Description

This function returns a vector of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on energy based effectiveness relative units. The BSWF is formulated for the range 210 nm to 400 nm.

### Usage

```
ICNIRP_e_fun(w.length)
```

### Arguments

w.length            numeric array of wavelengths (nm)

### Value

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (270 nm) and based on energy effectiveness.

### See Also

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [PG\\_q\\_fun\(\)](#)

## Examples

```
ICNIRP_e_fun(210:400)
```

---

IR

*Constructors of infra-red wavebands*

---

## Description

Wavelength-range definitions for *infrared* radiation according to ISO, CIE or as commonly defined in remote sensing applications.

## Usage

```
IR(std = "ISO")
```

```
NIR(std = "ISO")
```

```
IRA(std = "CIE")
```

```
SWIR(std = "CIE")
```

```
IRB(std = "CIE")
```

```
SWIR1(std = "RS")
```

```
SWIR2(std = "RS")
```

```
MIR(std = "ISO")
```

```
IRC(std = "CIE")
```

```
FIR(std = "ISO")
```

```
TIR1(std = "RS")
```

```
TIR2(std = "RS")
```

## Arguments

`std` character string, "ISO", "CIE", "RS" or Landsat imagers "LandsatRBV", "LandsatMSS", "LandsatTIRS", "LandsatOLI", "LandsatTM", "LandsatETM", depending on the constructor.

## Details

The values for `std = "ISO"` are according to ISO 20473. The values for `std = "CIE"` are suggested values according to Wikipedia, and need verification.

The wavelength limits for remote sensing `std = "RS"` and for Landsat imagers have been obtained from R package 'RStools' and NASA and USGS documentation.

The names NIR, SWIR and TIR are abbreviations of near infra-red, short-wave infra-red and thermal infra-red, respectively. The naming conventions are different for "CIE" than "ISO" standards, and the labels of the waveband objects reflect this with "IRA", "IRB", etc., used when appropriate.

## Value

A waveband object defining a wavelength range.

## See Also

[Far\\_red](#) for wavebands close to the boundary between red and infrared regions.

[new\\_waveband](#) and [waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

## Examples

```
SWIR1()  
SWIR1("RS")  
IR()  
NIR()  
MIR()  
IRA()  
IRB()
```

---

IR\_bands

*Constructor of lists of infrared wavebands*

---

## Description

Defined according to "ISO" or "CIE".

## Usage

```
IR_bands(std = "ISO")
```

## Arguments

`std` a character string "ISO" or "CIE".

**Value**

a list of wavebands

**See Also**

[waveband](#)

Other lists of unweighted wavebands: [Landsat\\_bands\(\)](#), [Plant\\_bands\(\)](#), [UV\\_bands\(\)](#), [VIS\\_bands\(\)](#)

**Examples**

```
IR_bands()
IR_bands("ISO")
IR_bands("CIE")
```

---

Landsat\_bands

*Constructor of lists of wavebands matching Landsat imagers*

---

**Description**

Defined according as ranges of wavelengths according to NASA and USGS manuals. The definitions are as *rectangular* windows, while the true response functions deviate to some extent from these ideal definitions.

**Usage**

```
Landsat_bands(std = "L8")

RBV_bands(std = "LandsatRBV")

MSS_bands(std = "LandsatMSS")

OLI_bands(std = "LandsatOLI")

TIRS_bands(std = "LandsatTIRS")

ETM_bands(std = "LandsatETM")
```

**Arguments**

`std` a character string "L1"..."L9", for missions, "LandsatRBV", "LandsatMSS", etc. for imagers.

**Details**

See <https://landsat.usgs.gov/spectral-characteristics-viewer> for detailed sensitivity spectra for the different bands of the imaginers.



**Value**

a list of wavebands

**See Also**

[waveband](#)

Other lists of unweighted wavebands: [IR\\_bands\(\)](#), [Plant\\_bands\(\)](#), [UV\\_bands\(\)](#), [VIS\\_bands\(\)](#)

**Examples**

```
Landsat_bands("L1")
Landsat_bands("L8")
OLI_bands()
TIRS_bands()
```

---

 NDVI

*Normalized Vegetation Index*


---

**Description**

Compute the NDVI from spectral reflectance according to waveband definitions from standards or corresponding to satellite imagers.

**Usage**

```
NDVI(spct, imager = "LandsatOLI", wb.trim = FALSE)
```

**Arguments**

spct	reflectance_spct or reflectance_mspct object.
imager	character Name of the imager or standard to be used.
wb.trim	logical Flag telling if wavebands crossing spectral data boundaries are trimmed or ignored.

**Details**

NDVI is used in remote sensing to the diagnose the condition of vegetation, including crops. It is used for Landsat imagery but also at the farm or plot scale using cameras on drones. It is computed as:

$$NDVI = (NIR - Red) / (NIR + Red)$$

The waveband ranges used to compute reflectance vary. Even the imagers in the different Landsat satellites 1 to 8 have had somehow different wavelength sensitivities. The `NDVI()` function uses the waveband constructors [Red](#) and [NIR](#) defined in this package. Reflectance is averaged over the wavebands using function [reflectance](#).

**Value**

A numeric vector. When the wavelength range of `spct` does not fully overlap with both wavebands NA is silently returned.

**Note**

The value passed as argument to `imager` must be a valid argument for both [Red](#) and [NIR](#). If the desired return value is a data frame, function [NDxI](#) can be used to flexibly compute NDVI and any similar index.

---

 Orange

---

*Constructor of orange waveband*


---

**Description**

Wavelength-range definition for *orange* radiation according to ISO.

**Usage**

```
Orange(std = "ISO")
```

**Arguments**

`std` a character string "ISO"

**Details**

Orange radiation (591...610 nm) as defined in ISO standards based on human colour vision.

**Value**

A waveband object defining a wavelength range.

**References**

ISO (2007) Space environment (natural and artificial) - Process for determining solar irradiances. ISO Standard 21348. ISO, Geneva.

**See Also**

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

## Examples

```
Orange()
Orange("ISO")

e_irrad(sun.spct, Orange()) # W m-2
q_irrad(sun.spct, Orange()) # mol m-2
q_irrad(sun.spct, Orange(), scale.factor = 1e6) # umol m-2
```

---

 PAR

*Constructors of PAR and PhR wavebands*


---

## Description

Waveband definitions for *photosynthetic radiation (PhR)*, *photosynthetically active radiation (PAR)* and *extended photosynthetically active radiation (ePAR)* according to different definitions in use for land plants.

## Usage

```
PAR(std = "PAR", norm = 550)
```

```
PhR()
```

## Arguments

std	a character string "Plant" (or "range"), "McCree" (or "photon", "PAR"), "Zhen" (or "ePAR"), "Gabielsen" (or "Gaastra" or "energy") or "Nichiporovich".
norm	normalization wavelength (nm)

## Details

Photosynthetically active radiation (400-700 nm) as proposed by McCree (1972), and currently used in plant sciences, gives equal weight to photons within its range, thus weights increase with increasing wavelength when expressed as energy. PAR is normally expressed as photon irradiance or (photosynthetic photon flux density, PPF) using implicitly 1 as weight for all wavelengths (a BSWF). It is also possible, but very unusual, to express the quantity PAR as defined in McCree (1972) as an *energy* irradiance, in which case a BSWF with weights different from 1 needs to be used. In this case the default normalization wavelength for the PAR BSWF is set at 550 nm (my own choice).

A proposal (see Zhen et al., 2021), defines extended photosynthetically active radiation (400-750 nm) as an alternative to PAR. The need to consider far-red photons as drivers of photosynthesis has become apparent with the increased use of LEDs for plant cultivation. **WARNING:** the proposed definition limits photon irradiance in the range 700-750 nm to a maximum of 30 larger than 1.4 times PAR even in the presence of far-red photons in excess, because far-red photons have only a synergistic effect on photosynthesis in PAR. Ensuring this condition is fulfilled is the responsibility

of the user of PAR(), ePAR() and PhR() functions. Currently, ePAR can usefully complement PAR, but in my opinion, it is far too early for it to replace PAR.

Some earlier definitions, described by McCree (1972a) citing Gabrielsen and Gaastra, used this same wavelength range but assuming wavelength-invariant response to energy within this same range, thus weights decrease with increasing wavelength when expressed as photons. McCree (1972a) cites Nichiporovich for a similar energy based quantity but covering a wider range of wavelengths (380-710 nm). McCree's definition from 1972 is currently the one preferred by most researchers and used almost universally in the plant sciences, while others are only of historical interest. Photosynthetic radiation (400-700 nm) is defined as a wavelength range and does not implement the spectral weighting inherent to McCree's (1972) definition of PAR or the earlier energy-based ones described by McCree (1972a).

### Value

For PhR, a waveband object defining a wavelength range . For PAR, a waveband object implementing the response curve of PAR as defined by McCree (1972) and thus including a weighting function used in computation of energy-base PAR irradiances. The weights are normalized to 1 at 550 nm. The waveband label is set to "PAR" or "PhR" accordingly.

### Warnings

PAR is sometimes described as a range of wavelengths (e.g., Both et al., 2015), which can be confusing as there is more to McCree's (1972) definition, an spectral response function by which all photons within the range of PAR elicit the same strength of response. As long as PAR is expressed as a photon irradiance or a photon irradiation, this, of course, makes no difference.

ePAR and PAR are meant to be use to quantify light sources with a broad spectrum, i.e., sources giving out white light or pale-coloured light. PAR and ePAR are technical measures of light useful for plants in the same way as illuminance is a measure of how bright light feels to an average human. They were never meant to describe the response to be expected from an individual in particular, be it a plant or human. They are generalizations, that allow us to consistently measure light in different situations.

### Note

PAR() and PhR() call the same function definition with different default arguments.

The default for the normalization wavelength at 550 nm keeps the average weights across the waveband equal to unity, except the special case of ePAR, where the photons in the extension to the range act by synergy.

Standard DIN 5031-10:2018-03 Defines two BSWFs *sy1* and *sy2* for photosynthesis, none of which are currently implemented in 'photobiologyWavebands'.

### References

- McCree, K. J. (1972) The action spectrum, absorptance and quantum yield of photosynthesis in crop plants. *Agricultural Meteorology*, 9, 191-216. doi:10.1016/00021571(71)900227
- McCree, K. J. (1972a) Test of current definitions of photosynthetically active radiation against leaf photosynthesis data. *Agricultural Meteorology*, 10, 443-453. doi:10.1016/00021571(72)900453

Both, A. J.; Benjamin, L.; Franklin, J.; Holroyd, G.; Incoll, L. D.; Lefsrud, M. G. & Pitkin, G. (2015) Guidelines for measuring and reporting environmental parameters for experiments in greenhouses. *Plant Methods*, 11:43. doi:10.1186/s1300701500835.

DIN (2018) Standard DIN 5031-10:2018-03 Optical radiation physics and illuminating engineering. Part 10: Photobiologically effective radiation, quantities, symbols and action spectra. Beuth Verlag, Berlin 2018

### See Also

[waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PG\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

### Examples

```
PAR()
PhR()
PAR("Plant")

q_irrad(sun.spct, PhR(), scale.factor = 1e6) # umol m-2 s-2
q_irrad(sun.spct, PAR(), scale.factor = 1e6) # umol m-2 s-2
e_irrad(sun.spct, PAR("Gabrielsen")) # W m-2
e_irrad(sun.spct, PhR()) # W m-2
e_irrad(sun.spct, PAR()) # BE W m-2, normalized at 700 nm
```

---

PG	<i>Constructor of PG weighted waveband</i>
----	--

---

### Description

Plant growth BSWF of Flint and Caldwell

### Usage

```
PG(norm = 300, w.low = 275, w.high = 390)
```

### Arguments

norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

### Details

The mathematical formulation included by Flint and Caldwell (2003) as an appendix is used. While this formulation is consistently used, the range of wavelengths over which it has been applied has varied. We use the approach of the NSF UV network and extrapolate up to 390 nm. The widely used simulation program TUV uses, instead, 366 nm as the boundary, which makes a significant difference to the computed irradiance values in sunlight.

### Value

a waveband object wavelength defining wavelength range, weighting function and normalization wavelength.

### Note

In the original publication [1], the long-end wavelength boundary is not specified. The longest wavelength at which the plant response was measured is 366 nm. From the data there is no evidence that action would immediately drop to zero at longer wavelengths. We have used in earlier versions the same value as used by the 'NSF Polar Programs UV Monitoring Network' as described in <https://web.archive.org/web/20220130091146/http://uv.biospherical.com/Version2/description-Version2-Database3.html>.

We use 390 nm as default value for w.high, but make if possible for the user to set a different wavelength. To reproduce the output of the TUV simulation model [3] version 5.0 set w.high = 366. The effect on the RAF and doses of changing this wavelength boundary is substantial, as discussed by Micheletti et al. [2]. Consequently, the value used must be always reported to ensure reproducibility. For comparisons with previous reports one may need to recompute effective irradiances using matching wavelength boundaries on a case by case basis.

### References

- [1] Flint, S. and Caldwell M. M. (2003) A biological spectral weighting function for ozone depletion research with higher plants *Physiologia Plantarum*, 2003, 117, 137-144
- [2] Micheletti, M. I.; Piacentini, R. D. & Madronich, S. (2003) Sensitivity of Biologically Active UV Radiation to Stratospheric Ozone Changes: Effects of Action Spectrum Shape and Wavelength Range *Photochemistry and Photobiology*, 78, 456-461
- [3] <https://www2.acom.ucar.edu/modeling/tropospheric-ultraviolet-and-visible-tuv-radiation-model>

### See Also

[GEN\\_G](#) [GEN\\_T](#) [GEN\\_M](#) and [waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [UV\\_health\\_hazard\(\)](#), [erythema\(\)](#)

### Examples

`PG()`  
`PG(300)`

---

PG_q_fun	<i>Gives values for the Plant Growth BSWF as a function of wavelength</i>
----------	---

---

### Description

This function gives a set of numeric multipliers that can be used as a weight to calculate effective doses and irradiances. The returned values are on quantum based effectiveness relative units.

### Usage

```
PG_q_fun(w.length)
```

### Arguments

w.length          numeric array of wavelengths (nm)

### Value

a numeric array of the same length as w.length with values for the BSWF normalized as in the original source (300 nm)

### Note

We follow the original definition here for the equation, with no limitation to the wavelength range. However, be aware that in practice it is not used for long wavelengths (different limits between 366 nm and 400 nm have been used by different authors).

### See Also

Other BSWF functions: [CH4\\_e\\_fun\(\)](#), [CH4\\_q\\_fun\(\)](#), [CIE\\_e\\_fun\(\)](#), [CIE\\_q\\_fun\(\)](#), [DNA\\_GM\\_q\\_fun\(\)](#), [DNA\\_P\\_q\\_fun\(\)](#), [FLAV\\_q\\_fun\(\)](#), [GEN\\_G\\_q\\_fun\(\)](#), [GEN\\_M\\_q\\_fun\(\)](#), [GEN\\_T\\_q\\_fun\(\)](#), [ICNIRP\\_e\\_fun\(\)](#)

### Examples

```
PG_q_fun(293:400)
```

---

photopic\_sensitivity    *Photopic sensitivity of the human eye*

---

**Description**

Constant value used in the definition of Lumen 1 W is equal to 683 Lumen at 555 nm

**Usage**

photopic\_sensitivity

**Format**

A single numeric value

**Details**

A single numeric value

---

Plant\_bands                      *Constructor of lists of wavebands used in plant biology*

---

**Description**

Defined according to different authors.

**Usage**

Plant\_bands(std = "sensory20")

**Arguments**

std                      a character string "sensory", "sensory10", "sensory20", "Sellaro", "ISO", "CIE", "none" or "", where "ISO", "CIE" and "none" affect only the UV bands.

**Value**

a list of wavebands

**See Also**

[waveband](#)

Other lists of unweighted wavebands: [IR\\_bands\(\)](#), [Landsat\\_bands\(\)](#), [UV\\_bands\(\)](#), [VIS\\_bands\(\)](#)



## Examples

```
Plant_bands()  
Plant_bands("sensory")  
Plant_bands("sensory10")  
Plant_bands("sensory20")  
Plant_bands("ISO")  
Plant_bands("CIE")
```

---

Purple	<i>Constructor of purple waveband</i>
--------	---------------------------------------

---

## Description

Wavelength-range definition for *purple* radiation according to ISO or imagers in the Landsat satellites.

## Usage

```
Purple(std = "ISO")
```

## Arguments

`std` a character string "ISO", or Landsat imager "LandsatOLI".

## Details

Purple (or violet) wavelengths as defined by ISO standards based on human vision overlap the UVA band as defined by a separate ISO standard. In other contexts like plant photobiology purple is included under blue, while some overoptimistic LED sellers call LEDs emitting in the violet region ultraviolet LEDs. In addition to the ISO definition of purple, a purple channel from Landsat imagers is implemented.

## Value

A waveband object defining a wavelength range.

## See Also

[new\\_waveband](#) [waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

**Examples**

```

Purple()
Purple("ISO")

e_irrad(sun.spct, Purple()) # W m-2
q_irrad(sun.spct, Purple()) # mol m-2
q_irrad(sun.spct, Purple(), scale.factor = 1e6) # umol m-2

```

---

Red *Constructor of red waveband*

---

**Description**

Wavelength-range definitions for *red* light according to ISO or as commonly used in plant or remote sensing applications.

**Usage**

```
Red(std = "ISO")
```

**Arguments**

std	a character string, one of "ISO", "Smith10", "Smith20", "Inada", "Warrington", "Sellaro", "RS", "LandsatOLI", "LandsatTM", "LandsatETM", "LandsatMSS", and "LandsatRBV".
-----	--

**Details**

The different arguments passed to formal parameter `std` determine the range of wavelengths set as boundaries of the returned waveband object; "ISO" is an standardized definition based on human colour vision; "Smith10", "Smith20", "Inada", "Warrington", "Sellaro" and "Broad" are non-standard but used in plant sciences; "RS" is non-standard but frequently used in remote sensing; the remaining definitions are for the published wavelength sensitivity range of imagers (cameras) in the Landsat satellite missions.

In plant photobiology the definitions proposed by Prof. Harry Smith are the most widely used, specially to compute a red to far-red photon ratio relevant to phytochrome photoreceptors. However, other authors have used different definitions in their publications. "Smith10" (655-665 nm), "Smith20" (650-670 nm), "Inada" (600-700 nm), "Warrington" (625-675 nm), and "Sellaro" (620-680 nm).

**Value**

a waveband object defining a wavelength range.

**Note**

The bands are defined as square windows, these can be applied to spectral data to obtain the "true" values, but they do not simulate the sensitivity of broad-band sensors or the spectral transmittance of ionic filters. Some band-pass interference filters may have very sharp cut-in and cut-off, and their effect can be approximated by a square window, but filters based on light absorption will show gradual tails and bell-shaped wavelength-windows. The Landsat instruments have very steep cut-in and cut-off slopes and are well approximated.

**References**

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). Beyond the Visible: A handbook of best practice in plant UV photobiology (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

ISO (2007) Space environment (natural and artificial) - Process for determining solar irradiances. ISO Standard 21348. ISO, Geneva.

Murakami, K., Aiga I. (1994) Red/Far-red photon flux ratio used as an index number for morphological control of plant growth under artificial lighting conditions. Proc. Int. Symp. Artificial Lighting, Acta Horticulturae, 418, ISHS 1997.

Sellaro, R., Crepy, M., Trupkin, S. A., Karayekov, E., Buchovsky, A. S., Rossi, C., & Casal, J. J. (2010). Cryptochrome as a sensor of the blue/green ratio of natural radiation in Arabidopsis. Plant physiology, 154(1), 401-409. [doi:10.1104/pp.110.160820](https://doi.org/10.1104/pp.110.160820).

Smith, H. (1982) Light quality, photoperception and plant strategy. Annual Review of Plant Physiology, 33:481-518. [doi:10.1146/annurev.pp.33.060182.002405](https://doi.org/10.1146/annurev.pp.33.060182.002405)

**See Also**

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

**Examples**

```
Red()
Red("ISO")
Red("Smith")
Red("Sellaro")

e_irrad(sun.spct, Red()) # W m-2
q_irrad(sun.spct, Red()) # mol m-2
q_irrad(sun.spct, Red(), scale.factor = 1e6) # umol m-2
```

---

scotopic\_sensitivity    *Scotopic sensitivity of the human eye*

---

**Description**

Constant value for human vision under very weak illumination 1 W is equal to 1699 Lumen at 507 nm

**Usage**

scotopic\_sensitivity

**Format**

A single numeric value

**Details**

A single numeric value

---

SetlowTUV.spct    *Setlow's action spectrum for DNA damage*

---

**Description**

A dataset containing the wavelengths at a 0.1 nm interval. Tabulated values for Setlow's naked DNA damage action spectrum as used in the TUV model.

**Format**

A response.spct object with 1082 rows and 2 variables

**Details**

The variables are as follows:

- w.length (nm)
- s.e.response

**References**

<https://web.archive.org/web/20220130091146/http://uv.biospherical.com/Version2/description-Version2.html> downloaded 2015-02-07

---

UV *Constructor of ultraviolet waveband*

---

**Description**

Wavelength-range definition for *ultraviolet (UV)* radiation according to ISO and CIE standards.

**Usage**

```
UV(std = "ISO")
```

**Arguments**

std	"ISO" or "CIE"
-----	----------------

**Details**

UV: 100–400 nm. The ranges agree between CIE and ISO standards, thus, the argument passed to parameter `std` only affects the labels in the returned waveband object.

**Value**

A waveband object defining a wavelength range.

**References**

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

**See Also**

[new\\_waveband waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

**Examples**

```
UV()  
UV("ISO")
```

---

UVA

*Constructor of ultraviolet-A waveband*

---

### Description

Wavelength-range definitions for *ultraviolet-A (UV-A)* radiation, by default according to ISO or as commonly used in different application areas.

### Usage

```
UVA(std = "ISO")
```

```
UVA1(std = "CIE")
```

```
UVA2(std = "CIE")
```

```
UVAsw(std = "plants")
```

```
UVA1w(std = "plants")
```

```
UVAsw(std = "plants")
```

### Arguments

std                    a character string "CIE", "ISO", "none", "medical" or "plants".

### Details

Implemented definitions. UV-A according to CIE and ISO standards: 315-400 nm. UV-A according to common non-standard practice: 320-400 nm. UV-A2 according to CIE report 134/1: 315-340 nm. UV-A1 according to CIE report 134/1: 340-400 nm. UV-Asw according to non-standard use possibly suitable for plants: 315-350 nm. UV-A1w according to non-standard use possibly suitable for plants: 350-400 nm.

### Value

A waveband object defining a wavelength range.

### Note

The non-standard definitions of UV-A and UV-A2 using 320 nm as limit should not be used in any new publications or work as they deviate from the internationally recommended practice. Their continued use leads to confusion. Their inclusion in this package is to allow calculations needed to compare new results and methods against old publications. UV-A1 and UV-A2 definitions are in wide use in medicine, but not yet standardized. Recent research on the plant photoreceptor UVR8 suggests that UV-A1 and UV-A2 bands are also relevant to plants (Rai et al., 2021). UV-A1w and UV-Asw have been used for plants, but UV-A1 and UV-A2 seem now preferable.

## References

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

CIE (1999) 134/1 TC 6-26 report: Standardization of the Terms UV-A1, UV-A2 and UV-B. <https://cie.co.at/publications/cie-collection-photobiology-photochemistry-1999>

Rai N, Morales LO, Aphalo PJ (2021) Perception of solar UV radiation by plants: photoreceptors and mechanisms. *Plant Physiology* 186: 1382–1396. [doi:10.1093/plphys/kiab162](https://doi.org/10.1093/plphys/kiab162)

## See Also

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

## Examples

```
UVA()
UVA("none")
UVA("ISO")
UVA("CIE")
```

```
UVA1()
UVA1("CIE")
```

```
UVA2()
UVA2("CIE")
```

```
e_irrad(sun.spct, UVA()) # W m-2
e_irrad(sun.spct, UVA1()) # W m-2
e_irrad(sun.spct, UVA2()) # W m-2
```

---

UVB

*Constructor of ultraviolet-B waveband*

---

## Description

Wavelength-range definitions for *ultraviolet-B*, *UV-B* radiation, by default according to ISO or as commonly used in different application areas.

## Usage

```
UVB(std = "ISO")
```

**Arguments**

std                    a character string "CIE", "ISO", "medical" or "none"

**Details**

Implemented definitions. UV-B according to CIE and ISO standards: 280–315 nm. UV-B according to common non-standard practice: 280–320 nm. UV-B according to medical or dermatological non-standard practice: 280–320 nm.

**Value**

a waveband object defining a wavelength range.

**Note**

The non-standard definition of UV-B using 320 nm as limit should not be used in any new publications or work as it deviates from the internationally accepted standards and its use leads to confusion. Its inclusion in this package is to allow calculations needed to compare new results and methods against old publications.

**References**

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

**See Also**

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

**Examples**

```
UVB()
UVB("ISO")
UVB("CIE")
UVB("none")
UVB("medical")

e_irrad(sun.spct, UVB()) # W m-2
q_irrad(sun.spct, UVB()) # mol m-2
q_irrad(sun.spct, UVB(), scale.factor = 1e6) # umol m-2
```



---

UVC                                      *Constructor of ultraviolet-C waveband*

---

### Description

Wavelength-range definitions for *ultraviolet-C (UV-C)* radiation, by default according to ISO or as commonly used in different application areas.

### Usage

```
UVC(std = "ISO")
```

### Arguments

`std`                                      a character string "CIE", "ISO", "none", or "medical".

### Details

Implemented definitions. UV-C according to CIE and ISO standards: 100–280 nm. UV-c according to common non-standard practice: 200–280 nm. UV-C according to medical or dermatological non-standard practice, e.g. Diffey (1991): 200–290 nm.

### Value

a waveband object wavelength defining a wavelength range.

### References

Aphalo, P. J., Albert, A., Björn, L. O., McLeod, A. R., Robson, T. M., Rosenqvist, E. (Eds.). (2012). *Beyond the Visible: A handbook of best practice in plant UV photobiology* (1st ed., p. xxx + 174). Helsinki: University of Helsinki, Department of Biosciences, Division of Plant Biology. ISBN 978-952-10-8363-1 (PDF), 978-952-10-8362-4 (paperback). Open access PDF download available at [doi:10.31885/9789521083631](https://doi.org/10.31885/9789521083631).

### See Also

[new\\_waveband waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UV\(\)](#), [VIS\(\)](#), [Yellow\(\)](#)

### Examples

```
UVC()  
UVC("CIE")  
UVC("ISO")  
UVC("none")  
UVC("medical")
```

---

UVI

*Calculate UV Index (UVI) from spectral irradiance*

---

### Description

UVI (UV Index) is a unitless quantity based on erythema biological spectral weighting function (BSWF), that gives an easy to interpret UV measure, mainly meant for informing general public about sunburn risk.

### Usage

```
UVI(spct, std = "NOAA", integer_UVI = FALSE)
```

### Arguments

spct	a source.spct object
std	character One of "WMO", "NOAA".
integer_UVI	logical Return a positive integer value according to WMO recommended practice or a numeric value.

### Details

Two different definitions of UV Index are implemented in this package. Setting `std="NOAA"` follows the definition in Kiedron et al. (2007) but using CIE98 as SWF. NOAA definition discards wavelengths shorter than 286.5 nm as when calculated based on spectral data from Brewer instruments. "WMO" uses the internationally accepted lower limit at 250 nm (see WHO, 2002). "NOAA" is the default as this is safer with noisy data for solar radiation measured at ground level, and in this case the value of UVI should be correct, and almost identical except for errors caused by noise at shorter wavelengths. However, when calculating UVI from radiation spectra from UV lamps, "WMO" should be used, as most UV lamps do emit some radiation between 250 nm and 286.5 nm.

### Value

depending on the argument passed to `integer_UVI`, a numeric (FALSE) or integer (TRUE) value for the unitless UVI.

### Note

The default is to return a numeric value not rounded into a value in the integer-number based recommended UVI scale. This is done to avoid loss of precision in cases when additional operations, such as averaging, are applied to the UVI values.

## References

World Health Organization, World Meteorological Organization, United Nations Environment Programme & International Commission on Non-Ionizing Radiation Protection. (2002) Global Solar UV Index: A Practical Guide. World Health Organization, Geneva. ISBN 9241590076. <https://apps.who.int/iris/handle/10665/42459>

P. Kiedron, S. Stierle and K. Lantz (2007) Instantaneous UV Index and Daily UV Dose Calculations. NOAA-EPA Brewer Network. <https://www.esrl.noaa.gov/gmd/grad/neubrew/docs/UVindex.pdf>

## Examples

```
UVI(sun.spct)
UVI(sun.spct, "WMO")
UVI(sun.spct, integer_UVI = TRUE)
```

---

UV\_bands

*Constructor of lists of UV wavebands*

---

## Description

Defined according to "ISO" by default, but other definitions also supported.

## Usage

```
UV_bands(std = "ISO")
```

## Arguments

std                    a character string "ISO", "CIE", "medical", "plants" or "none".

## Value

a list of wavebands

## See Also

[waveband](#)

Other lists of unweighted wavebands: [IR\\_bands\(\)](#), [Landsat\\_bands\(\)](#), [Plant\\_bands\(\)](#), [VIS\\_bands\(\)](#)

## Examples

```
UV_bands()
UV_bands("ISO")
UV_bands("CIE")
UV_bands("medical")
UV_bands("plants")
UV_bands("none")
```

---

UV\_health\_hazard      *Constructor of UV health hazard weighted waveband*

---

### Description

Waveband constructor for ICNIRP UV health hazard 2004 BSWF.

### Usage

```
UV_health_hazard(std = "ICNIRP", norm = 270, w.low = 210, w.high = 400)
```

```
ICNIRP(norm = 270, w.low = 210, w.high = 400)
```

### Arguments

std	a character string "ICNIRP".
norm	normalization wavelength (nm)
w.low	short-end boundary wavelength (nm)
w.high	long-end boundary wavelength (nm)

### Details

This BSWF is used for the determination of exposure limits (EL) for workers, and includes a safety margin as it is based on eye and the non-pathologic response of the most sensitive human skin types when not tanned. Values are interpolated according to equations 2a, 2b and 2c in ICNIRP (2004), which cover the range 210 nm to 400 nm.

**The program code is provided as is, with no guarantee of suitability for any purpose, and should under no circumstances be used to assess actual health hazards.**

### Value

a waveband object defining wavelength range, weighting function and normalization wavelength.

### Note

The weights at 180, 190, 200 and 205 nm are presented only in tabular in ICNIRP (2004) and all values at wavelengths < 210 nm taken as NA.

Standard DIN 5031-10:2018-03 defines BSWF *uvh* as a table of interpolated values derived from ICNIRP UV health hazard. So, the values computed using this R package do not necessarily exactly match those according to DIN 5031-10:2018-03. The range of wavelengths used here, 210 to 400 nm, does not agree, with those in the standard: 180 to 400 nm.

## References

INTERNATIONAL COMMISSION ON NON-IONIZING RADIATION PROTECTION (2004) ICNIRP GUIDELINES ON LIMITS OF EXPOSURE TO ULTRAVIOLET RADIATION OF WAVELENGTHS BETWEEN 180 nm AND 400 nm (INCOHERENT OPTICAL RADIATION). HEALTH PHYSICS 87(2):171-186. <https://www.icnirp.org/cms/upload/publications/ICNIRPUV2004.pdf>

## See Also

[new\\_waveband](#) [waveband](#)

Other BSWF weighted wavebands: [CH4\(\)](#), [DNA\\_GM\(\)](#), [DNA\\_N\(\)](#), [DNA\\_P\(\)](#), [FLAV\(\)](#), [GEN\\_G\(\)](#), [GEN\\_M\(\)](#), [GEN\\_T\(\)](#), [PAR\(\)](#), [PG\(\)](#), [erythema\(\)](#)

## Examples

```
ICNIRP()
UV_health_hazard()
UV_health_hazard("ICNIRP")
```

---

VIS

*Constructor of VIS waveband*

---

## Description

Visible (to humnas) radiation (380...760 nm) according to ISO standard definition, no weighting applied. For `std = "RS"` the returned range is the same as for `PAR()`. The panchromatic bands of Landsat missions are also supported.

## Usage

```
VIS(std = "ISO")
```

## Arguments

`std` a character string "ISO" or "RS" (remote sensing).

## Value

A waveband object wavelength defining a wavelength range.

## See Also

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [Yellow\(\)](#)

**Examples**

```
VIS()  
VIS("ISO")  
VIS("LandsatOLI")  
VIS("Landsat7")  
VIS("Pan.RBV.Landsat3")
```

---

VIS\_bands

*Constructor of lists of VIS wavebands*

---

**Description**

Defined according to "ISO".

**Usage**

```
VIS_bands(std = "ISO")
```

**Arguments**

std                    a character string "ISO".

**Value**

a list of wavebands

**See Also**

Other lists of unweighted wavebands: [IR\\_bands\(\)](#), [Landsat\\_bands\(\)](#), [Plant\\_bands\(\)](#), [UV\\_bands\(\)](#)

**Examples**

```
VIS_bands()  
VIS_bands("ISO")
```

---

Yellow	<i>Constructor of yellow waveband</i>
--------	---------------------------------------

---

**Description**

Wavelength-range definition for *yellow* radiation according to ISO.

**Usage**

```
Yellow(std = "ISO")
```

**Arguments**

std                    a character string "ISO"

**Details**

Yellow radiation (570...591 nm) as defined in ISO standards based on human colour vision.

**Value**

a waveband object wavelength defining a wavelength range.

**See Also**

[waveband](#)

Other unweighted wavebands: [Blue\(\)](#), [Far\\_red\(\)](#), [Green\(\)](#), [IR\(\)](#), [Orange\(\)](#), [Purple\(\)](#), [Red\(\)](#), [UVA\(\)](#), [UVB\(\)](#), [UVC\(\)](#), [UV\(\)](#), [VIS\(\)](#)

**Examples**

```
Yellow()
Yellow("ISO")

e_irrad(sun.spct, Yellow()) # W m-2
q_irrad(sun.spct, Yellow()) # mol m-2
q_irrad(sun.spct, Yellow(), scale.factor = 1e6) # umol m-2
```

# Index

## \* BSWF functions

CH4\_e\_fun, 8  
CH4\_q\_fun, 9  
CIE\_e\_fun, 11  
CIE\_q\_fun, 12  
DNA\_GM\_q\_fun, 13  
DNA\_P\_q\_fun, 16  
FLAV\_q\_fun, 21  
GEN\_G\_q\_fun, 23  
GEN\_M\_q\_fun, 25  
GEN\_T\_q\_fun, 27  
ICNIRP\_e\_fun, 29  
PG\_q\_fun, 39

## \* BSWF weighted wavebands

CH4, 7  
DNA\_GM, 13  
DNA\_N, 14  
DNA\_P, 16  
erythema, 17  
FLAV, 20  
GEN\_G, 22  
GEN\_M, 24  
GEN\_T, 26  
PAR, 35  
PG, 37  
UV\_health\_hazard, 52

## \* datasets

CIE1924\_lef.spct, 9  
CIE1951\_scotopic\_lef.spct, 10  
CIE2008\_lef2deg.spct, 11  
photopic\_sensitivity, 40  
scotopic\_sensitivity, 44  
SetlowTUV.spct, 44

## \* lists of unweighted wavebands

IR\_bands, 31  
Landsat\_bands, 32  
Plant\_bands, 40  
UV\_bands, 51  
VIS\_bands, 54

## \* unweighted wavebands

Blue, 5  
Far\_red, 18  
Green, 28  
IR, 30  
Orange, 34  
Purple, 41  
Red, 42  
UV, 45  
UVA, 46  
UVB, 47  
UVC, 49  
VIS, 53  
Yellow, 55

Blue, 5, 20, 29, 31, 34, 41, 43, 45, 47–49, 53, 55

CH4, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53  
CH4\_e\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39

CH4\_q\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39

CIE (erythema), 17

CIE1924\_lef.spct, 9

CIE1951\_scotopic\_lef.spct, 10

CIE2008\_lef2deg.spct, 11

CIE\_e\_fun, 8, 9, 11, 12, 14, 17, 21, 24, 26, 27, 29, 39

CIE\_q\_fun, 8, 9, 12, 12, 14, 17, 21, 24, 26, 27, 29, 39

DNA\_GM, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53

DNA\_GM\_q\_fun, 8, 9, 12, 13, 17, 21, 24, 26, 27, 29, 39

DNA\_N, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53

DNA\_N\_q\_fun, 15



- DNA\_P, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53
- DNA\_P\_q\_fun, 8, 9, 12, 14, 16, 21, 24, 26, 27, 29, 39
- erythema, 7, 13, 14, 16, 17, 21, 22, 25, 26, 37, 38, 53
- ETM\_bands (Landsat\_bands), 32
- Far\_red, 6, 18, 29, 31, 34, 41, 43, 45, 47–49, 53, 55
- FIR (IR), 30
- FLAV, 7, 13, 14, 16, 18, 20, 22, 25, 26, 37, 38, 53
- FLAV\_q\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39
- GEN.G, 26
- GEN.M, 26
- GEN\_G, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53
- GEN\_G\_q\_fun, 8, 9, 12, 14, 17, 21, 23, 26, 27, 29, 39
- GEN\_M, 7, 13, 14, 16, 18, 21, 22, 24, 26, 37, 38, 53
- GEN\_M\_q\_fun, 8, 9, 12, 14, 17, 21, 24, 25, 27, 29, 39
- GEN\_T, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 53
- GEN\_T\_q\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39
- Green, 6, 20, 28, 31, 34, 41, 43, 45, 47–49, 53, 55
- ICNIRP (UV\_health\_hazard), 52
- ICNIRP\_e\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39
- IR, 6, 20, 29, 30, 34, 41, 43, 45, 47–49, 53, 55
- IR\_bands, 31, 33, 40, 51, 54
- IRA (IR), 30
- IRB (IR), 30
- IRC (IR), 30
- Landsat\_bands, 32, 32, 40, 51, 54
- MIR (IR), 30
- MSS\_bands (Landsat\_bands), 32
- NDVI, 33
- NDxI, 34
- new\_waveband, 6, 13, 14, 16, 25, 31, 41, 45, 49, 53
- NIR, 20, 33, 34
- NIR (IR), 30
- OLI\_bands (Landsat\_bands), 32
- Orange, 6, 20, 29, 31, 34, 41, 43, 45, 47–49, 53, 55
- PAR, 7, 13, 14, 16, 18, 21, 22, 25, 26, 35, 38, 53
- PG, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 37, 53
- PG\_q\_fun, 8, 9, 12, 14, 17, 21, 24, 26, 27, 29, 39
- photobiologyWavebands  
(photobiologyWavebands-package), 3
- photobiologyWavebands-package, 3
- photopic\_sensitivity, 40
- PhR (PAR), 35
- Plant\_bands, 32, 33, 40, 51, 54
- Purple, 6, 20, 29, 31, 34, 41, 43, 45, 47–49, 53, 55
- RBV\_bands (Landsat\_bands), 32
- Red, 6, 20, 29, 31, 33, 34, 41, 42, 45, 47–49, 53, 55
- reflectance, 33
- scotopic\_sensitivity, 44
- SetlowTUV.spct, 44
- SWIR (IR), 30
- SWIR1 (IR), 30
- SWIR2 (IR), 30
- TIR1 (IR), 30
- TIR2 (IR), 30
- TIRS\_bands (Landsat\_bands), 32
- UV, 6, 20, 29, 31, 34, 41, 43, 45, 47–49, 53, 55
- UV\_bands, 32, 33, 40, 51, 54
- UV\_health\_hazard, 7, 13, 14, 16, 18, 21, 22, 25, 26, 37, 38, 52
- UVA, 6, 20, 29, 31, 34, 41, 43, 45, 46, 48, 49, 53, 55
- UVA1 (UVA), 46
- UVA2 (UVA), 46
- UVA1w (UVA), 46
- UVAsw (UVA), 46
- UVB, 6, 20, 29, 31, 34, 41, 43, 45, 47, 47, 49, 53, 55

UVC, [6](#), [20](#), [29](#), [31](#), [34](#), [41](#), [43](#), [45](#), [47](#), [48](#), [49](#),  
[53](#), [55](#)

UVI, [50](#)

VIS, [6](#), [20](#), [29](#), [31](#), [34](#), [41](#), [43](#), [45](#), [47–49](#), [53](#), [55](#)

VIS\_bands, [32](#), [33](#), [40](#), [51](#), [54](#)

waveband, [6](#), [7](#), [13](#), [14](#), [16](#), [18](#), [20–22](#), [25](#), [26](#),  
[29](#), [31–33](#), [37](#), [38](#), [40](#), [41](#), [43](#), [45](#),  
[47–49](#), [51](#), [53](#), [55](#)

Yellow, [6](#), [20](#), [29](#), [31](#), [34](#), [41](#), [43](#), [45](#), [47–49](#),  
[53](#), [55](#)